

Formation of Radial Electric Field Shear at Boundary of Magnetic Island in LHD

M.Yoshinuma, K. Ida, K.Tanaka, R. Sakamoto, M. Hoshino, T. Morisakim, A. Komori
and the LHD Experimental Group

National Institute for Fusion Science, Toki, Japan

e-mail contact of main author: yoshinuma@nifs.ac.jp

Abstract. Active control of a radial electric field shear by a combination of pellet injection and a magnetic island induced by external perturbation coils is demonstrated in the Large Helical Device (LHD). In this experiment, the radial electric field shear at the boundary of the magnetic island reaches 0.3 MV/m^2 and is sustained by repetitive injection of the pellet and the improvement of ion transport is observed near the boundary of the magnetic island.

1. Introduction

Since the radial electric field, E_r , and its shear has been recognized to play an important role in the improvement of transport, the effects of the E_r and its shear on the transport have been investigated in various magnetic confinement devices. Active control of the radial profile of the radial electric field is one of the important tools to investigate the effects of the E_r and its shear. In low temperature plasmas, biased electrodes experiments have been performed to investigate effects of radial electric field and/or its shear on fluctuations and anomalous transport in linear devices [1-3], in torus devices [4] in order to change radial electric field actively. There is a difficulty to insert the electrode into plasmas to control the radial electric field because of the high heat flux to electrode from the plasma in high temperature plasmas. Therefore there is few biased electrodes experiment in large toroidal confinement devices where the plasma temperature is too high to insert the electrode. In tokamak plasmas, the transition of the radial electric field and the formation of radial electric field shear are observed associated with the transition from L-mode to H-mode plasma [5, 6]. In helical plasmas, the radial electric field is expected to reduce the neoclassical ripple loss in low collisional regime. The reduction of the ion thermal diffusivity due to the large positive electric field is observed associated with the transition from small E_r to large positive E_r in the LHD when the neoclassical ion loss is dominant loss [7]. On the other hand, the strong E_r shear in the plasma with an internal transport barrier (ITB) contributes to the reduction of turbulent transport as observed in the compact helical system (CHS), Wendelstein-7AS and the LHD [8-12] by the local heating with center focused ECH. In the plasma with the internal transport barrier in the LHD, the reduction of electron thermal diffusivity is observed associated with the appearance of the E_r shear near the boundary between the electron root in the core and ion root in the edge with the combination of center focused ECH and NBI. Thus, the effect of E_r shear in the electron root is well investigated in helical systems related with the ITB. The effect of the E_r shear in ion root on thermal diffusivity has not been much investigated. We demonstrate an active control of the radial electric field shear between large negative E_r in the core and small E_r in the edge in the ion root regime by a combination of pellet injection and a non-rotating magnetic island induced by external perturbation coils in LHD.

2. Scheme of E_r Shear formation in an Ion root plasma

In general, the radial electric field determined by the neo-classical non-ambipolar ion and electron flux can be controlled by changing the magnitude of helical ripples [13] and collisionality [14] in helical plasmas. The radial electric field becomes positive (in




FIG. 1. Schematic view of poloidal cross section of magnetic flux surface near the magnetic island induced by external perturbation coil and the E_r profile by pellet injection.

the electron root) when the plasma collisionality is low enough, while it becomes negative (in the ion root) at higher collisionality. The collisionality can be increased large enough to make the plasma to be in the ion root by increasing the electron density using gas puffing or pellet injection. Since the radial electric field tends to be zero inside the magnetic island [15], the large radial electric field shear will be produced at the boundary of the magnetic island in the ion root regime where the E_r is negative outside of the magnetic island. Figure 1 show the schematic view of this experiment. The radial electric field shear near the boundary of a $n/m=1/1$ non-rotating magnetic island induced by external perturbation coils (Local Island Divertor coils: LID coils) is expected to be produced and sustained with continuous fueling by the repetitive pellet injections.

3. Target discharge and pellet injection for E_r formation

The experiment was performed in an NBI sustained plasma with a magnetic axis of 3.6m, averaged minor radius of 0.6m and the magnetic field strength of 2.5T on the LHD. Figure 2 (a) shows heating scheme by the NBI and ECH which used to start up of the plasma. Six pellets with a volume of 12 mm³ are injected with a repetition time of 0.15 sec at the timing indicated by arrow in Fig. 2(b) to produce negative E_r in the ion root plasma. The electron density at the plasma center and the stored energy is increased up to $3.7 \times 10^{19} \text{ m}^{-3}$ and 400 kJ at $t=1.86\text{sec}$ associated with the repetitive pellet injection as shown in Fig. 2(b) and (c). The deposition power of NBI, which is depend on the density, is 4.2 MW at $t=1.86\text{sec}$. The radial electric field is derived from the poloidal rotation velocity at the mid-plane of the vertically elongated cross section with the charge exchange spectroscopy (CXS) [14,16] using the charge exchange line of fully ionized Neon. The Neon impurity is injected by gas puff before the pellet injection to make CXS available during the formation of E_r as shown in Fig. 2(b). Figure 2(d) shows the time evolution of the radial electric field at $R=4.04\text{m}$ measured with the charge exchange spectroscopy with the time-resolution of 150msec. The increase of the electron density is small enough to keep the plasma to stay in the electron root with the Neon puff. The radial electric field changes from positive (electron root) to negative (ion root) associated with increasing the electron density with repetitive injections of pellet. The critical density of the transition between positive and negative is about $1.5 \times 10^{19} \text{ m}^{-3}$. The strong negative E_r (ion root) is achieved at $t=1.4 \text{ sec}$ by the repetitive pellet injection and sustained up to the end of the NBI. Figure 3 shows the time




FIG.2. Time trace of (a) port through power of NBI and the timing of ECH, (b) line averaged electron density and timing of gas puffing and pellet injection, (c) stored energy, (d) radial electric field at $\rho=0.75$.




FIG.3. Time evolution of the radial electric field at the $R=4.04$ with repetitive (circle), multi (square) and single pellet injection (triangle).

evolution of the radial electric field with three type of pellet injection. Although the negative E_r is formed near the magnetic island just after the pellet injection at $t=1.05$ sec, the formation of negative electric field is transient and the negative E_r changes back to positive in the time scale of few 100 msec in the discharge with a single pellet injection. In the discharges with multi pellet injection (inject at $t=1.05, 1.2, 1.35$ sec) and repetitive pellet injection, the negative electric field can be sustained even after the last injection of the pellet. The volume of the pellet used for single injection and multi injection is 21 mm^3 which is larger than that used for the repetitive injections (12 mm^3). Faster change of E_r is observed in the discharge with larger pellet injection. It is considered




FIG4. Time traces of line averaged electron density in the plasmas both with (solid line) and without (dashed line) magnetic island at high density and with magnetic island (dotted line) at low density.

that the time scale of the E_r formation is related to the amount of the particle fuelling.

4 Required condition of E_r shear formation in an Ion root plasma

Figure 4 shows the three time traces of the line averaged electron density in the plasma both with and without magnetic island at high density and with magnetic island at low density. The line averaged electron density is increased up to over the critical density in the high density discharge while the line averaged electron density is lower than the critical density in the low density discharge. The effect of magnetic island on particle transport is relatively small because the two time traces of the line averaged density at high density discharge are almost identical to each others.

Figure 5 shows the radial profiles of E_r after the pellet injection ($t=1.7s$) measured by charge exchange spectroscopy with time resolution of 400 msec which can be measured with wide range in space compared with the CXS system with high-time resolution (150msec). The radial electric field is already close to zero in the plasma near the critical density and therefore there is no E_r shear at the boundary of magnetic island in the low density discharge. In the plasma with the electron density higher than the critical density, the small negative E_r is observed at the inner region of the plasma ($\rho < 0.74$) and its magnitude gradually increases toward the plasma edge when there is no magnetic island. On the other hand, when there is a magnetic island, the large negative E_r is observed except for the inside of the magnetic island, where the E_r vanishes because of damping of the poloidal flow, and large E_r shears are observed at the boundaries of the magnetic island.

Figure 6 shows the radial profiles of the ratio of electron density 0.1 sec after ($t=1.15$ sec) the start of the pellet injection ($t=1.05$ sec) to that 0.1 sec before ($t=0.95$ sec) the start of pellet injection in the plasma with and without the magnetic island. Although the ablation location indicated by the peak of H_α intensity is at $r=0.68-0.86$, the peak of increase in density appears at $r=0.82-0.94$ because the ablation cloud moves outward due to $E_\theta \times B$ force. The larger increase of the density is observed in the plasma with the magnetic island as compared with it in the plasma without the magnetic island and the more negative E_r is produced in the plasma with the magnetic island as compared with it in the plasma without the magnetic island at the inner region of the plasma. These observations clearly show that both the magnetic island and high density plasma by pellet injection are necessary for formation of the E_r shear in an ion root.

4. Time evolution of E_r shear and gradient of ion pressure




FIG.5. Radial profiles of E_r after the pellet injection in the plasmas with (closed circles) and without (open squares) magnetic island at high density and with magnetic island (open triangles) at low density.




FIG.6. The radial profile of the ratio of electron density 0.1 sec after the pellet injection to that 0.1 sec before the pellet injection ($t=1.05\text{sec}$) in the plasma with and without the magnetic Island.

Figure 7 shows the time evolution of the radial profile of E_r in the plasma with a magnetic island. Where the Δt indicates the time difference from the start of the pellet injection ($t=1.05\text{sec}$). The E_r changes from positive ($\Delta t=-0.8\text{s}$) to negative ($\Delta t > 0.22\text{s}$) on the inner side of the magnetic island while there are no large changes in the E_r inside the magnetic island as the line averaged electron density increases from $0.7 \times 10^{19} \text{m}^{-3}$ to $2.3\text{-}3.2 \times 10^{19} \text{m}^{-3}$ by repetitive pellet injections. The radial electric field shear (dE_r/dR) also changes its sign from negative to positive associated with the change of the E_r sign at $\rho=0.79$. Thus, the E_r shear can be controlled by changing the collisionality outside of the magnetic island in helical plasmas. Figure 8 shows the time evolutions of the E_r shear (dE_r/dR) at $\rho=0.79$. With the repetitive pellet injection, the positive E_r shear becomes strong and reaches 0.3MV/m^2 0.3sec after the first pellet injection in the plasma with the magnetic island while the no significant E_r shear is observed in the plasma without the magnetic island. This large E_r shear can be sustained during the repetitive pellet injection with a repetition time of 0.15sec. Because the electron density is lower than the critical density before the pellet injection, the E_r inner side of the magnetic island is positive (electron root) and the E_r shear is negative. When the electron density exceeds the critical density by the pellet injection, the E_r is negative




Fig.7: Radial profiles of the E_r 0.08s before and 0.07, 0.22, 0.52s after the pellet injection.

(ion root) and the E_r shear is positive after the pellet injection.

Figure 9 shows the time evolution of ion pressure gradient $-n_e dT_i/dR$ at $\rho=0.77$. The gradient of ion temperature, dT_i/dR , is derived from the ion temperature profiles measured with the CXS, while the electron density, n_e , is measured with an FIR interferometer. As shown in Fig.9, the value of the $-n_e dT_i/dR$ in the plasma with the magnetic island is almost twice that in the plasma without the magnetic island associated with the positive E_r shear formation. This experiment demonstrates that the ion transport is improved associated with an appearance of negative E_r shear. On the other hand, there is no sign of transport improvement due to the positive E_r shear as seen in the ion pressure at $t=0.8-1.05$. It is noted that there is no difference in ion heat flux in these plasmas because the electron density and NBI power are controlled to be identical. The ion thermal diffusivity χ_i is estimated to be reduced by a factor of two near the magnetic island within a few cm by creating the large positive E_r shear at the boundary of magnetic island with the assist of pellet injection in ion root plasma. In the ion root, the increase of the ion pressure gradient makes E_r to be more negative. It is considered that there is feed back between the E_r shear, reduction of χ_i , and the increase of the ion pressure gradient. When the E_r is negative, the feed back process becomes positive feedback because the larger ion pressure due to the reduction of χ_i contributes more negative E_r and larger E_r shear as shown in Fig.10. On the other hand, when the E_r is positive, it becomes negative feedback and no improvement of the ion transport is expected.

5. Summary

Large E_r shear in the ion root regime can be produced and sustained at the boundary of the magnetic island by combination of repetitive pellet injection and a magnetic island induced by external perturbation coils. The electron density which is much higher than the critical density is required to produce the E_r shear in an ion root. The gradient of ion pressure is increased up to twice of that in the plasma without the magnetic island associated with the formation of the positive E_r shear in the ion root plasma, while there is no differences of the ion pressure gradient between with and without the magnetic island in the negative E_r shear in the electron root. It was suggested that there is a positive feedback process between the E_r shear, reduction of χ_i , and increase of ion pressure gradient when the E_r is negative (ion root) and dE_r/dR is positive. It may be that the positive feedback process contributes to sustaining the E_r shear. The combination of a magnetic island and pellet injection is considered to be a useful tool to control E_r shear and an ion energy transport near the magnetic island.




Fig.8: Time evolutions of the E_r shear in plasmas with (circle) and without (square) a magnetic island. Arrows indicate the timings of pellet injection.




Fig.9: Time evolutions of the ion pressure gradient at $\rho \sim 0.77$ in plasmas with (circle) and without (square) a magnetic island.

Acknowledgement

The authors would like to thank the LHD technical staff for their effort to support the experiments in LHD. They also acknowledge continuous encouragement and support of Director-General O. Motojima. This work is supported by NIFS05ULBB512 and partly supported by a grant-in-aid for scientific research of MEXT Japan.

References

- [1] A. Tsushima, et al, "Control of Radial Potential Profile and Nonambipolar Ion Transport in an Electron Cyclotron Resonance Mirror Plasma", *Phys.Rev.Let.* **56** (1986) 1815.
- [2] A. Komori, et al, "Control of low-frequency plasma instabilities by a nonuniform radial electric field", *Phys. Fluids* **31** (1988) 210.
- [3] M.Yoshinuma, et al., "Effects of Radial Electric Field and its Shear on Low-Frequency Fluctuations in the Q-T-U Device", *Trans. Fusion Tech.* **39** (2001)




FIG.10. Positive feed back between the E_r shear, reduction of ion diffusivity, and increase of ion pressure gradient.

- 191.
- [4] TAYLOR, R. J., et al., "H-Mode Behavior Induced by Cross-Field Currents in a Tokamak", Phys. Rev. Lett. **63** (1989) 2365.
- [5] GROEBNER, R. J., et al., "Role of edge electric field and poloidal rotation in the L-H transition", Phys. Rev. Lett. **64** (1990) 3015.
- [6] IDA, K., et al., "Edge electric-field profiles of H-mode plasmas in the JFT-2M tokamak", Phys. Rev. Lett. **65** (1990) 1364.
- [7] IDA, K., et al., "Reduction of Ion Thermal Diffusivity Associated with the Transition of the Radial Electric Field in Neutral-Beam-Heated Plasmas in the Large Helical Device", Phys. Rev. Lett. **86** (2001) 5297.
- [8] FUJISAWA, A., et al., "Electron Thermal Transport Barrier and Density Fluctuation Reduction in a Toroidal Helical Plasma", Phys. Rev. Lett. **82** (1999) 2669.
- [9] STROTH, U., et al., "Internal Transport Barrier Triggered by Neoclassical Transport in W7-AS", Phys. Rev. Lett. **86** (2001) 5910.
- [10] SHIMOZUMA, T., et al., "Formation of electron internal transport barriers by highly localized electron cyclotron resonance heating in the large helical device", Plasma Phys. Control. Fusion **45** (2003) 1183.
- [11] IDA, K., et al., "Characteristics of Electron Heat Transport of Plasma with an Electron Internal-Transport Barrier in the Large Helical Device", Phys. Rev. Lett. **91** (2003) 085003-1.
- [12] TAKEIRI, Y., et al., "Formation of electron internal transport barrier and achievement of high ion temperature in Large Helical Device", Phys. Plasma, **10** (2003) 1788.
- [13] IDA, K., et al., "Control of the radial electric field shear by modification of the magnetic field configuration in LHD", Nucl. Fusion, **45** (2005) 391.
- [14] YOSHINUMA, M., et al., "Observations of edge radial electric field transition in LHD plasmas", Plasma Phys. Control. Fusion, **46** (2004) 1021.
- [15] IDA, K., et al., "Observation of Plasma Flow at the Magnetic Island in the Large Helical Device", Phys. Rev. Lett., **88** (2002) 15002.
- [16] IDA, K., et al., "Measurements of poloidal rotation velocity using charge exchange spectroscopy in a large helical device", Rev. Sci. Instrum., **71** (2000) 2360.