

Studies of Stimulated Raman Scattering in Laser Plasma Interactions

A. Salcedo, R. J. Focia, A. K. Ram, and A. Bers

Plasma Science & Fusion Center, M.I.T., Cambridge, MA, 02139, U.S.A.

E-mail contact of main author: bers@mit.edu

Abstract. Coupled theoretical and computational work is presented aimed at understanding and modeling stimulated Raman backscattering (SRBS) relevant to laser-plasma interactions (LPIs) in large-scale, nearly homogeneous plasmas. We address the following five observed, nonlinear phenomena associated with SRBS: coupling of SRBS to Langmuir decay interactions (LDIs); cascading of LDI; SRS cascades; and stimulated electron acoustic wave scattering (SEAS).

1. Introduction

In the past few years, experiments on the TRIDENT facility at the Los Alamos National Laboratory (LANL) have undertaken to study LPIs in (independently) pre-formed plasmas with a diffraction limited laser beam, thus approximately simulating LPIs in a single hot spot (SHS) [1], relevant to indirect drive inertial confinement fusion. We present results from recent analytical and computational studies [2,3] of LPIs with particular reference to SRBS in SHS [3]–[5].




Figure 1: (a) $k\lambda_{De} = 0.32$ ($n_e/n_{cr} = 0.04$); (b) $k\lambda_{De} = 0.28$ ($n_e/n_{cr} = 0.05$).

2. Coupled Mode Model Equations (CMME) for SRBS–LDI and Cascades

We considered the following model LPI in their simplest description of nonlinearly coupled modes [6]: (a) SRBS; (b) SRBS coupled to LDI; (c) SRBS + LDI coupled to LDI-cascade (LDIc); and (d) SRBS coupled to SRS-cascade (SRSc). In the most complex case considered, (c), this entailed solving the following seven nonlinearly coupled mode equations:

$$\text{LASER} : \quad L_1 a_1 = -K_1 a_2 a_3 \quad (1)$$

$$\begin{aligned}
\text{BEMW(EPW)} &: L_2 a_2 = K_1 a_1 a_3 & (2) \\
\text{EPW(BEMW)} &: L_3 a_3 = K_1 a_1 a_2 - K_2 a_4 a_5 & (3) \\
\text{BEPW(FEMW)} &: L_4 a_4 = K_2 a_3 a_5 - K_3 a_6 a_7 & (4) \\
\text{IAW(EPWc)} &: L_5 a_5 = K_2 a_3 a_4 & (5) \\
\text{CEPW} &: L_6 a_6 = K_3 a_4 a_7 & (6) \\
\text{CIAW} &: L_7 a_7 = K_3 a_4 a_6 & (7)
\end{aligned}$$

In these equations $L_j = (\partial/\partial t) + v_{gj}(\partial/\partial x) + \nu_j$ is the linear wavepacket operator on the slowly-varying action density amplitudes in space and time (see first reference in [6]). The above equations can be reduced to apply to (d) by labeling (2) as EPW, (3) as BEMW, (4) as FEMW, (5) as EPWc [as shown in parentheses along (2)–(5)], and setting K_3 , a_6 , and a_7 to zero.




Figure 2: (a) SRBS reflectivity with and without coupling to LDI; (b) SRBS reflectivity with and without the SRS cascade.

2.1 Bounded Length Interaction Computations From CMME Studies

We modeled the SHS-LPI as occurring in a bounded homogeneous plasma of length L , with an incident laser field at $x = 0$. Parameters chosen to represent typical SHS-LPI experiments [1,5] were: $L = 250 \mu m$; a Maxwellian plasma with 70% hydrogen and 30% carbon ions; $T_e \approx 700 eV$; $T_i \approx 100\text{--}500 eV$ (depending on electron density n_e where the interaction was made to occur); $\lambda_o = 527 nm$; and a laser intensity $P_o \approx 10^{15} \text{ Watts/cm}^2$; (n_e/n_{cr}) was varied, as in the experiments to obtain LPIs as a function of the EPW $k\lambda_{De}$. The results shown in the following figures are presented in terms of normalized quantities used in the computations: action densities normalized to laser a_0 , and distances normalized to $v_{g3}/(|K_{LDI}|a_0)$. Thus for the given laser and plasma parameters, the interactions length of $250 \mu m$ is 900 normalized x -units, and the SRBS energy flow reflectivity is

$$SRBS_r \equiv \left. \frac{|v_{g2}|w_2}{|v_{g1}|w_1} \right|_{x=0} \approx \frac{\omega_2}{\omega_1} |a_2(x=0, t)|^2 . \quad (8)$$

(Note, $x = 0$ is shown as -450 normalized x units.) The following is a summary of results.

1. *Coupling of SRBS to LDI* — Dramatic changes in space-time occur as the ELD of the EPWs is changed. For strong ELD ($k\lambda_{De} \gtrsim 0.4$), although SRBS may




Figure 3: (a) Electron distribution at 51.7 (ω_{pe}^{-1}); (b) Its deviation from the initial Maxwellian.

be present, LDI is not excited anywhere in the SHS plasma. For intermediate ELD ($k\lambda_{De} \sim 0.32$), LDI is excited but spatially localized near the boundary where the laser enters the SHS plasma; the EPW is saturated just above the LDI threshold [see Figure 1(a)]. Notably, for weak ELD ($k\lambda_{De} \lesssim 0.28$), LDI is intense in the plasma region near the laser entrance boundary, and exhibits incoherent space-time fluctuations [Fig. 1(b)] akin to spatiotemporal chaos (STC) [7]; the consequence is an appreciable reduction in SRBS reflectivity due to the ensuing dephasing. However, the low reflectivities observed in the SHS-LPI experiments are not predicted by SRBS coupling to LDI [Figure 2(a)].

2. *Cascading of LDI* — The SHS-LPI experiments [3] have shown the existence of LDI cascades associated with SRBS. From our coupled mode simulations of (1)–(7), that include the first LDI cascade, we find that its effect is to slightly increase the SRBS reflectivity. This can be understood from the fact that the cascade drains energy from the LDI, thus enhancing SRBS which was reduced by LDI in the absence of the cascade. We thus explain the observation of cascades and their effect on the SRBS reflectivity, but we find that these cascades do not contribute significantly to predict the observed saturated values of reflectivity in the particular SHS-LPI.
3. *SRS Cascades* — We find that an SRS cascade gives a more significant decrease in the SRBS reflectivity than LDI [Fig. 2(b)]. This is readily understood from the Manley-Rowe relations. SHS-LPI experiments have so far not looked for this type of cascade.

Our results, although different in some respects, are complementary to [8,9].

3. Fluid Vlasov-Maxwell Simulations and Models for SRBS–SEAS

We have studied and modeled SEAS, analytically and numerically, with a model that differs from the one proposed in [4], considering that SRBS evolves its EPW to ampli-

tudes that trap electrons. With a nonlinear, one-dimensional, Eulerian, Vlasov-Maxwell code [10], we launched a laser propagating field on a trapped electron distribution function due to an independently generated EPW in a plasma with parameters of the recent SEAS experiment [3,4], thus simulating a saturated state of SRBS. The evolution of this setup shows indeed that SRBS is essentially saturated, and in addition exhibits the generation of low-density, thermally spread beamlets below the phase velocity of the EPW (Figure 3) [11]. Estimating from Fig. 3(b) the presence of a beamlet ($v_b \approx 0.045 c$, $T_b \approx 40 eV$, $n_b \approx 0.02 n_e$), the linear natural modes of the plasma with beamlet show a relatively weakly damped EAW [see Figure 4(a)]. Parametric growth rates for this




Figure 4: (a) EPW and EAW modes; (b) Nonlinear electrostatic modes [13].

system, driven by the laser, are given by the kinetic dispersion relation (see e.g., [12]):

$$(1 + \chi_e) D_+ D_- = (kv_0/2)^2 \chi_e (D_+ + D_-) \quad (9)$$

where $\chi_e = [-Z'(\zeta_p)/2k^2\lambda_{Dp} - Z'(\zeta_b)/2k^2\lambda_{Db}]$, with $\zeta_p = (\omega/|k|\sqrt{2}v_{Tp})$, and $\zeta_b = (\omega - kv_b/|k|\sqrt{2}v_{Tb})$; $D_{\pm} = c^2(k \pm k_0)^2 + \omega_{pe} - (\omega \pm \omega_0)^2$, and $v_0 = (e|E_0|/m_e\omega_0)$. For laser intensities of 10^{16} Watts/cm², calculations from (9) show that for a range of beamlet parameters (from $T_b \lesssim 50 eV$ for $n_b \gtrsim 0.02 n_e$, and $T_b \lesssim 20 eV$ for $n_b \gtrsim 0.01 n_e$), consistent with Fig. 3(b), one obtains a (quasimode) parametric growth rate $\approx 1.6 \times 10^{-4} \omega_o$ in a narrow frequency range at $\omega_r \approx 0.45 \omega_{pe}$ ($k\lambda_{De} \approx 0.285$), similar to the observed SEAS, and an SRBS growth rate of $\approx 9.4 \times 10^{-3} \omega_o$ (essentially unaffected by the above beamlet parameters) in a narrow range at $\omega_r \approx 1.15 \omega_{pe}$ ($k\lambda_{De} \approx 0.267$), as expected. Preliminary results from the (k, ω) spectra of the Vlasov code simulations confirm the results of this model. Finally, we find that a fully nonlinear, *steady state* solution of the one-dimensional Vlasov equation in an electrostatic wave electric field [13] exhibits *nonlinear* EAW-type modes below the plasma frequency [see Fig. 4(b), $E_N = eE_o/(m_e\omega_p\sqrt{2}v_{Te})$]; their excitation in SRBS also remains to be studied.

4. Acknowledgments

This work was supported in part by Los Alamos National Laboratory (LANL) Contract E29060017-8F, and in part by Department of Energy Contract DE-FG02-91ER-

54109 and National Science Foundation/Department of Energy Contract DE-FG02-99ER-54555. We are grateful to Dr. Magdi Shoucri of the Institut de Recherche de l'Hydro Quebec (IREQ) for making his Vlasov code available to us and for help in carrying out the simulations, and to Mr. David Strozzi of M.I.T. for computations of the kinetic, parametric growth rates.

5. References and Footnotes

- [1] MONTGOMERY, D. S., et al., "Characterization of Plasma and Laser Conditions for Single Hot Spot Experiments," *Laser and Particle Beams* **17** (1999) 349.
- [2] SALCEDO, A., Coupled Modes Analysis of SRS Backscattering, with Langmuir Decay and Possible Cascadings, Ph.D. Thesis, EECS, M.I.T. (January 2002).
- [3] FOCIA, R. J., Investigation and Characterization of Single Hot Spot Laser-Plasma Interactions, Ph.D. Thesis, EECS, M.I.T. (January 2002).
- [4] MONTGOMERY, D. S., et al., "Observation of Stimulated Electron-Acoustic-Wave Scattering," *Phys. Rev. Lett.* **87** (2001) 155001.
- [5] FERNANDEZ, J. C., et al., "Current Status of Laser-Matter Interaction Experimental Research at Los Alamos National Laboratory," presented 26th European Conf. Laser Interaction with Matter (ECLIM) Prague, 2000.
- [6] BERS, A., "Linear Waves and Instabilities," *Plasma Physics – Les Houches 1972* (DEWITT, C., PEYRAUD, J., Eds.), Gordon and Breach Publishers, New York (1975) 113–215; HEIKKINEN, J. A., et al., "Intensity Saturation of Stimulated Raman Scattering by Ion-Wave Coupling," *Phys. Fluids* **29** (1986) 1291; BONNAUD, G., PESME, D., "Stimulated Raman Scattering and Ion Dynamics: The Role of Langmuir Wave Non-Linearities," *Laser Interaction & Related Plasma Phenomena* (Proc. 8th Int. Workshop Monterey, California, 1987), Vol. 8, Plenum Press, New York (1988) 375–399.
- [7] RAM, A. K., et al., "Saturation of SRS by Spatiotemporal Chaos in Coupled Langmuir Decay," *Controlled Fusion & Plasma Physics 1994* (Proc. 21st EPS Montpellier, France, 1994), (JOFFRIN, E., et. al., Eds.), Vol. III, EPS, Geneva (1994) 1460–1463.
- [8] BONNAUD, G., et al., "Nonlinear Raman Scattering Behavior With Langmuir and Sound Waves Coupling in a Homogeneous Plasma," *Phys. Fluids B* **2** (1990) 1618.
- [9] KOLBER, T., et al., "Saturation of Stimulated Raman Scattering by Langmuir and Ion-Acoustic Wave Coupling," *Phys. Fluids B* **5** (1993) 138.
- [10] BERTRAND, P., et al., "A Nonperiodic Euler-Vlasov Code for the Numerical Simulation of Laser-Plasma Beat Wave Acceleration and Raman Scattering", *Phys. Fluids B* **2** (1990) 1028.
- [11] Because of limitations in the current state of the code, it was not possible to ascertain the persistence of the beamlets for times longer than the transit time ($\approx 170 \omega_p^{-1}$) of the laser through the simulated length ($\approx 80 \mu m$) — albeit this time is some 4–5 SRBS growth times. Thus the long-time persistence of the beamlets has not been established so far with the code, and they may well be a transient phenomenon. However, in the experimental setup [1,3], the SHS region is continuously replenished with fresh plasma so that, on the average, the beamlets and their effects appear in a quasi steady state. For the present we assume this to be the case.
- [12] DRAKE, J. F., et al., "Parametric Instabilities of Electromagnetic Waves in Plasmas," *Phys. Fluids* **17** (1974) 778.
- [13] KRAPCHEV, V., RAM, A. K., "Adiabatic Theory for a Single Nonlinear Wave in a Vlasov Plasma," *Phys. Rev. A* **22** (1980) 1229.